




**"ANTI-HYPERTENSIVE COMPOSITION
AND METHODS OF TREATMENT"**

I CLAIM:

ELST AVAILABLE COPY

1. A pharmaceutical composition for the treatment of hypertension comprising an effective anti-hypertensive amount of at least one compound in association with a pharmaceutically acceptable, substantially non-toxic carrier or excipient; said compound having one of the formulae:



wherein: R and R' may be the same or different and are H, alkyl or aralkyl having from 1 to 12 carbon atoms;
R₁ - R₆ may be the same or different and are H, R or R';
R₇ is H, alkyl, aryl or aralkyl having from 1 to 12 carbon atoms;

FILED AVAILABLE ON

m and n may be the same or different and
are integers from 3 to 10, inclusive;
a - e may be the same or different and
are integers from 3 to 10, inclusive;

or

(IV) a salt of (I), (II) or (III) with a pharmaceutically
acceptable acid.

2. A composition according to claim 1 wherein m is
3 and n is 4.

3. A composition according to claim 1 wherein m
and n are 3.

4. A composition according to claim 1 wherein m
and n are 4.

5. A composition according to claim 1 wherein R
and R' are alkyl.

6. A composition according to claim 1 wherein R
and R' are aralkyl.




7. A composition according to claim 1 wherein R
and R' are methyl.

8. A composition according to claim 1 wherein R and R' are ethyl.

9. A composition according to claim 1 wherein R and R' are propyl.

10. A composition according to claim 1 wherein R₁ and R₆ are benzyl.

11. A method of effecting anti-hypertensive action which comprises administering to a patient requiring anti-hypertensive effect at least one compound having one of the formulae:



wherein: R and R' may be the same or different and are H, alkyl or aralkyl having from 1 to 12 carbon atoms;
R₁ - R₆ may be the same or different and are H, R or R';

R₇ is H, alkyl, aryl or aralkyl having
from 1 to 12 carbon atoms;
m and n may be the same or different and
are integers from 3 to 10, inclusive;
a - e may be the same or different and
are integers from 3 to 10, inclusive;

or

(IV) a salt of (I), (II) or (III) with a pharmaceutically
acceptable acid.

12. A method according to claim 11 wherein m is 3
and n is 4.

13. A method according to claim 11 wherein m and n
are 3.

14. A method according to claim 11 wherein m and n
are 4.

15. A method according to claim 11 wherein R and R'
are alkyl.

16. A method according to claim 11 wherein R and R'
are aralkyl.

17. A method according to claim 11 wherein R and R'
are methyl.

1 18. A method according to claim 11 wherein R and R'
2 are ethyl.

1 19. A method according to claim 11 wherein R and R'
2 are propyl.

1 20. A method according to claim 11 wherein R and R'
2 are benzyl.